



OMRI
Listed

BIOMIN[®] CALCIUM

Chelated Mineral for Soil and Foliar Application



PRODUCT INFORMATION

BIOMIN CALCIUM contains chelated calcium designed for soil and foliar application. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct micronutrient deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizer.

DIRECTIONS AND

**GENERAL RECOMMENDATIONS:
SHAKE WELL BEFORE USE**

SOIL APPLICATION: BIOMIN CALCIUM can be applied before or at planting as banding or broadcast. The application rate is 1 quart-1 gallon per acre (2.5 to 10 liters/hectare) depending on soil conditions and application methods. For maintenance, 1 quart per acre (2.5 liters/hectare) is recommended. **BIOMIN CALCIUM** should be diluted with enough water to cover the entire acre.

FOLIAR APPLICATION:

FIELD CROPS: Apply 1-3 quarts per acre (2-8 liters/hectare), when crops are about 6-12 inches (15 - 30 cm) high or when deficiency symptoms appear. Repeat the application in 2 - 4 weeks if necessary.

FRUIT TREES AND GRAPES: Apply 1-3 quarts per acre (2-8 liters/hectare) starting at bud break or when deficiency symptoms ap-

GUARANTEED ANALYSIS

Calcium (Ca).....5.0%
5.0% chelated calcium
Derived from calcium carbonate, citric acid (chelating agent).

ALSO CONTAINS NONPLANT FOOD INGREDIENT: 10% Hydrolyzed vegetable protein.

CAUTION- Do not apply this product after application of copper containing products as this may cause phytotoxicity.

PREVENTS AND CORRECTS CALCIUM DEFICIENCIES

KEEP OUT OF REACH OF CHILDREN

MANUFACTURED BY:

JH BIOTECH, INC.
P.O. Box 3538
Ventura, CA 93006, U.S.A

NET CONTENTS:

- 1 Pint (0.47 liters)
- 2.5 Gallons (9.45 liters)
- 2 x 2.5 Gallons (18.9 liters)
- 5 Gallons (18.9 liters)
- 55 Gallons (207.9 liters)
- Other: _____ Gallons

Weight per Gallon: 9.0 lbs. (4.1 kg.)
@ 68°F

U.S. Pat.# 5,504,055

LOT NO.:

pear. Repeat the application in about 4 weeks if necessary.

VEGETABLE CROPS: Apply 1-3 quarts per acre (2-8 liters/hectare) at about 2 - 4 weeks after emergence, or when deficiency symptoms appear. Repeat the application in 1 - 3 weeks if necessary.

For best results, apply when the crop is in active growing stage, after irrigation or natural rain fall. For foliar application, it should be diluted with at least 100 times of water. **BIOMIN CALCIUM** will lower high pH of water and decrease the hydrolysis of most pesticides.

BIOMIN CALCIUM can be applied to strawberry, lettuce, tomato, celery, grapes, stone fruits, tree fruits, vegetables, commercial turf, and ornamental gardens.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

JH Biotech, Inc.

0301-CA-A-03
Copyright © 2011 JH Biotech, Inc.
All rights reserved.